

journal homepage: www.elsevier.com/locate/csbj

Mini Review

Comparative genomics for biodiversity conservation

Catherine E. Grueber*

Faculty of Veterinary Science, University of Sydney, NSW, Australia
San Diego Zoo Global, San Diego, CA, USA

ARTICLE INFO

Article history:

Received 29 January 2015

Received in revised form 13 May 2015

Accepted 15 May 2015

Available online 21 May 2015

Keywords:

Adaptation

Disease

Hybridisation

Management units

Phylogeny

Threatened species

ABSTRACT

Genomic approaches are gathering momentum in biology and emerging opportunities lie in the creative use of comparative molecular methods for revealing the processes that influence diversity of wildlife. However, few comparative genomic studies are performed with explicit and specific objectives to aid conservation of wild populations. Here I provide a brief overview of comparative genomic approaches that offer specific benefits to biodiversity conservation. Because conservation examples are few, I draw on research from other areas to demonstrate how comparing genomic data across taxa may be used to inform the characterisation of conservation units and studies of hybridisation, as well as studies that provide conservation outcomes from a better understanding of the drivers of divergence. A comparative approach can also provide valuable insight into the threatening processes that impact rare species, such as emerging diseases and their management in conservation. In addition to these opportunities, I note areas where additional research is warranted. Overall, comparing and contrasting the genomic composition of threatened and other species provide several useful tools for helping to preserve the molecular biodiversity of the global ecosystem.

© 2015 Grueber. Published by Elsevier B.V. on behalf of the Research Network of Computational and Structural Biotechnology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction	370
1.1. Applications of comparative genomics to conservation	371
1.1.1. Characterisation of conservation units	371
1.1.2. Informing the conservation consequences of hybridisation	371
1.2. Discovering the drivers of divergence	372
1.3. Understanding intrinsic threats and disease resilience	373
2. Conclusions	373
Acknowledgments	373
References	373

1. Introduction

Conservation genetics has entered the world of genomics [1]. The number of species with whole-genome sequence data is continually growing [2,3], so that more and more endangered taxa are becoming “genome-enabled” [4], that is, genome resources are available for them or their close relatives. These new technologies provide researchers with unprecedented levels of data to generate precise estimates of essential population genetic parameters, to examine questions

such as the causes and genetic consequences of population decline and fragmentation [1,5; for critique see 6]. Most applied conservation genetics research targets issues operating within or amongst populations of the same species (which may be spatially or temporally separated) [7]. This level of focus is often appropriate because anthropogenic threatening processes typically occur over relatively short evolutionary time frames: the scale relevant to population/species-level processes rather than deeper evolutionary trajectories such as speciation. Nevertheless, there is additional insight to be gained from considering the evolutionary context of threatened species, i.e. by taking a comparative approach across taxa. For example, comparative analyses of species' demographic and life history characteristics have revealed those particular ecological traits that predispose species to high risk of extinction [8,9].

* Faculty of Veterinary Science, University of Sydney, NSW, Australia.
E-mail address: catherine.grueber@sydney.edu.au.

In this paper, I explore how comparative approaches using genomic data may also add value to conservation efforts.

Comparative genomics benefits most from high-quality, annotated and mapped genome data, but the pre-existence of such complete data is not necessarily a prerequisite for taking a comparative genomic approach to wildlife genetics [10]. This is good news for conservation scientists, who frequently work on non-model species for which genome resources do not exist. Several options are available, although not all will be uniformly applicable across contexts: reduced-representation libraries [11] provide a cost-effective means of obtaining genome-level data for comparative studies [12]. RNAseq to obtain transcriptome-level data can also provide valuable insight, without necessarily obtaining whole-genome data (e.g. comparative RNA sequencing of 12 primate species, most of which had little or no genomic resources [13]). Genome-informed SNP arrays, developed for well-studied species, can be used to generate large amounts of data for closely related threatened taxa (e.g. utilising a primarily domestic dog SNP array to study wild canids [14]), although the percentage of shared polymorphisms between species decays exponentially with divergence time, decreasing the amount of data obtained from the chip for more distant species [15]. A further approach to preliminary comparative genomics investigation is the generation of large amounts of sequencing data, which is then aligned to the annotated genome of a closely related species (e.g. aligning California condor sequencing data against the chicken genome [16]).

Wildlife genomics may be undertaken at multiple levels, from comparing individuals within a population (in a population genetics/genomics framework) to comparisons at higher taxonomic levels (comparative genomics). Many genomics techniques offer opportunities for conservation (for recent overviews, see [1,3,6,17]). However, despite their potential value, comparative genomic studies with explicit and specific conservation applications remain uncommon ([1,6,18], exceptions are [19,20]). Impediments to the uptake of genomics in conservation include sampling and analysis constraints [21], as well as a lack of clear examples of successful application [6], amongst others. In this review, I focus on possible applications of comparative genomics to conservation, and provide examples of a variety of avenues for future work in this field. Comparative genomics itself is a broad field, with the potential to answer many salient questions in evolutionary biology, medicine, and other fields (e.g. [22]), and therefore the analyses mentioned herein also have many applications beyond threatened species management. In fact, due to the scarcity of conservation examples, much of the empirical work I discuss here has been conducted on non-threatened species. I touch on a number of topics in brief: my aim is not to provide an exhaustive survey, but rather an overview of new ways that an ever-growing resource of genomic data can be exploited to address timely problems in biodiversity conservation.

1.1. Applications of comparative genomics to conservation

My main discussion centres on a summary of conservation science research questions that may be approached or supported by the use of comparative genomic methods, and identification of research needs to further progress these aims.

1.1.1. Characterisation of conservation units

Identifying units of conservation is a fundamental goal of any conservation strategy, essential to both resource planning in a legal and financial sense (e.g. how to distribute conservation effort) and management planning in a practical and biological sense (e.g. which populations may be mixed and which show important distinctiveness that should be preserved). Although definitions vary [23], the concept of conservation management units encompasses groupings beyond traditional taxonomic demarcations, such as evolutionarily significant units and/or variants with particular ecological or social value.

Nevertheless, conservation units are usually informed by phylogeny, traditionally using putatively neutral genetic regions such as microsatellite markers or mtDNA. Importantly, these methods inform conservationists as to the degree of migration amongst putative conservation units [24] providing a distinction between “evolutionarily significant units”: populations that are phylogenetically discrete, and “management units”: populations with significant divergence in allele frequencies [25].

Recently, researchers have begun to target adaptive molecular variation for inclusion in the assessment of conservation units. These data introduce information about evolutionary distinctiveness into the definition of protected populations [26]. For example, diversity and differentiation at the major-histocompatibility complex (MHC), genes associated with adaptive immunity [27], have been incorporated into the delineation of conservation management units for several species, such as giant panda *Ailuropoda melanoleuca* [28] and marbled murrelets *Brachyramphus marmoratus* (a threatened seabird) [29]. However, basing management decisions on a small number of functional genomic regions presents a high risk of failing to detect evolutionarily and ecologically important processes that influence other parts of the genome [30]. Recent studies have shown how genome-level data can provide very high resolution for the reconstruction of phylogenetic trees, enabling detailed identification of species boundaries and relationships [12,31]. For example, Wagner et al. [31] recently used reduced-representation RAD sequencing to generate exceptionally detailed phylogenetic inference amongst 16 cichlid species in Lake Victoria, a community well-studied in evolutionary ecology. Evolutionary relationships amongst these species had previously been difficult to dissect using traditional methods, due to very recent divergence times which impaired discrimination amongst morphologically distinct species using much smaller numbers of nuclear and mitochondrial DNA sequence variants [31].

In conservation, phylogenetic approaches have been used to identify the most evolutionarily distinct species, which may then be targeted for particular conservation effort (e.g. EDGE [evolutionarily distinct, globally endangered] species [32]). Taking a whole-genome comparative approach to the characterisation of conservation units provides at least three advantages over traditional approaches: 1) greater resolution via the use of many more loci, 2) the ability to incorporate a wide diversity of putatively functional genetic regions (i.e. genic sequences) and 3) the ability to perform analyses using either neutral or functional data (or both), enabling researchers to study how different processes drive population structure [33]. Several challenges exist with the use of whole-genome data for the reconstruction of phylogenetic trees, such as how to conduct inference regarding species trees in the case of conflicting gene trees from different genomic regions [34]. These issues apply to all studies that use multigene data for phylogenetics, not just those with conservation aims, and their resolution is still an area of active research (e.g. [34,35]). Nevertheless, genome-level data enables researchers to determine whether any differentiation observed amongst populations results from evolutionary or demographic processes. For example, genetic structures based on different genomic regions (such as microsatellites versus MHC) are frequently uncorrelated (e.g. [29]), typically interpreted as a greater role of selection than drift at immunogenetic versus neutral loci, respectively [36]. Populations differing as a result of recent, drift-associated processes are not considered as distinct as populations differing as a result of deep adaptation processes [23]. Differentiating these mechanisms of structure amongst populations is essential to the fully informed preservation and management of molecular biodiversity in ecosystems [1].

1.1.2. Informing the conservation consequences of hybridisation

Human landscape modification has increased the frequency with which hybridisation influences the evolutionary course of many species around the world [37]. Introgression of a threatened species by a previously geographically separated and more-common relative can affect species integrity and result in extinction of the rarer type [38,39].

Although the term “hybridisation” typically refers to interspecific breeding, it has also been used to refer to interbreeding at lower levels of genetic differentiation, which may be of conservation relevance, such as amongst sub-species or regional variants. Hybridisation between previously separated groups may not only result in the extinction of rare forms by assimilation, but also accelerate the decline of threatened populations via outbreeding depression: the decreased fitness that occurs when distantly related lineages interbreed, as a result of the break-up of locally adapted haplotypes or gene combinations [40–42]. Determining the evolutionary timing of hybridisation is important from a conservation perspective, in order to evaluate whether it has occurred as a result of natural or anthropogenic processes [39]. Population and comparative genetics can be used to quantify the degree of ancient or recent hybridisation between threatened and non-threatened species [14,43], as well as to identify taxon-specific markers for the diagnosis and monitoring of hybridisation in dynamic systems (e.g. [44]). Identifying the degree of hybridisation represented by particular individuals, and thus whether any “pure forms” remain in a population, is essential in a conservation context for determining whether pure lineages can be recovered [39].

Studying hybridisation in some taxonomic groups requires molecular markers at the genome-level due to peculiar characteristics of the species’ genomic architecture (such as a recent [~25–100 MYA] genome duplication in salmonids [20], or polyploidization in plants [45]). In addition, research at the comparative genomic level can provide additional information that cannot be obtained using conventional genetic approaches: by examining extensive functional diversity data, researchers can identify the particular genomic regions that determine the physiological consequences of hybridisation. For example, diagnosing particular chromosomal regions under strong selection during hybridisation events allowed researchers to quantify the proportion of the wheat genome that has been affected by introgression [45]. Such studies can also inform whether particular genes are experiencing strong selection as a result (e.g. the identification of “super invasive” alleles at higher-than-expected frequencies in admixed populations of salmonids [46]).

Identifying genomic regions most affected by introgression may also provide diagnostic markers for rapidly distinguishing hybrids from pure forms [46]. These approaches have been used to inform the management of westslope cutthroat trout *Oncorhynchus clarki lewisi* in North America [20,46], for which the major threat to persistence is hybridisation with introduced rainbow trout *O. mykiss* [47]. Researchers used genome-level data to identify 3180 species-diagnostic SNPs, which were then genotyped in fish from multiple populations to evaluate levels of introgression and numbers of pure forms remaining [20,46]. The data also provided estimates of the proportion of the genomes of individual hybrid fish that could be traced back to cutthroat or rainbow trout. Quantifying individual-level introgression in this way enabled the researchers to study introgression with great precision, including identification of those animals for which a very small proportion of their genome came from the invasive species, as well as the discovery of candidate adaptive regions experiencing strong selection pressure during the hybridisation process [46].

Overall, the comparative genomic approach can provide unprecedented levels of precision for the study of hybridisation in conservation contexts, providing diagnostic markers for the identification of hybrids versus pure forms. Comparative genome-level analyses have the potential to reveal the genomic architecture of hybridisation, for the purpose of understanding the evolutionary mechanisms that drive hybrid genome evolution. From a practical perspective, these findings can also predict the effectiveness of novel selective breeding approaches to producing pure forms from recently introgressed populations as well as identifying those individuals to target (e.g. [48,49]). Molecular selective breeding may enable the recovery of genetic diversity contained within hybrids, which would otherwise be lost if only pure forms were targeted for breeding [49]. Using molecular genetics for

the latter is preferred over pedigree or morphological approaches to preserving rare forms from hybridisation, which can result in very high levels of inbreeding [50], although there are advantages to integrating multiple data types, especially when there may be ascertainment bias in the reference genomic material relative to the target species (e.g. [51]). The use of genome-level molecular data for de-introgression is still in its infancy: so far it has been pursued theoretically using real molecular data from intentional hybrids of Merino and Poll Dorset sheep breeds [49], and proposed in conservation planning for the endangered Cika cattle breed, which has been historically hybridised with other breeds [51]. Computational modelling has shown that selective breeding regimes could ultimately recover pure genomes from introgressed populations in only a few generations [49]. It will be interesting to observe whether this approach, currently targeting the conservation of rare breeds of commercially important domestic species, will offer benefits for the recovery of wildlife species threatened by hybridisation in the future.

1.2. Discovering the drivers of divergence

A broad goal of comparative genomics is to reveal the evolutionary forces that drive genetic and genomic diversity and, as already shown, these findings can add value to specific conservation problems. Importantly, comparative approaches can reveal the genomic regions under selection and putative genetic underpinnings of unique traits [3,13,52]. Identifying regions of the genome that are more divergent than expected, as well as those regions that are more variable than expected, may help guide the preservation of genetic diversity in conservation management, especially in captivity. At present, little is known about the ongoing genetic consequences of adaptation to captivity in threatened species, particularly when captive-raised animals are reintroduced to the natural environment [23]. Comparing genomes amongst species can help to identify those regions that are experiencing particularly strong positive selection, which may signify rapid adaptation to captivity, providing markers for the management of this undesired process [23,53].

At the whole-genome level, comparative approaches can reveal gene losses/gains and help researchers discover the defining characteristics of species (e.g. [54,55]) as these changes are major contributors to functional evolution and divergence [56]. However, inference based on gene losses/gains requires very high-quality genomes [57]. Many conservation genomic studies use data obtained through reduced-representation methods or other unassembled data types [58], which would not be suitable for addressing these questions. It has also been noted that such “genome scan” population genomic studies need to be undertaken with caution, as the risk of false-positives is high [59]. Continued advances in the computational comparison of genomes, in order to more accurately catalogue regions of divergence between species (whether sequence divergence or gene gains/losses) may provide valuable information about the biology and life history of species (e.g. [54]). Identifying the genetic origin of unique traits can be helpful for managing threatening processes (e.g. determining whether they have intrinsic or extrinsic causes). For example, in comparing the giant panda genome to those of other carnivores, the absence of digestive cellulose enzymes enabled Li et al. [55] to infer that the strict bamboo diet of the species is unlikely to have arisen as a result of intrinsic genetic characteristics, and may be directed more by gut microbiota. In addition, identifying regions of shared synteny between related species can aid in the characterisation of highly divergent orthologous genes, which would be difficult to identify using homology searches alone [54]. Clearly, there are advantages to be gained from taking a comparative approach to discovering the most divergent loci, and a better understanding threatened species and their potential response to threats. Ongoing improvements to genomic assembly and comparison, particularly the identification of variation in gene composition amongst species, will help to facilitate this conservation research.

1.3. Understanding intrinsic threats and disease resilience

Over time, small, isolated populations will gradually lose genetic diversity and levels of inbreeding will increase; together, these processes erode the resilience of these populations to environmental changes and lead to increased expression of deleterious recessive alleles [60,61]. These genetic impacts leave populations vulnerable to a variety of novel threats, particularly diseases. Comparative genomic approaches can assist in the study of the disease threats that arise from losses in genetic diversity, by providing a greater understanding of the origin of diseases and the response of populations. The results can then be used to design strategies for disease treatment or management. In this section, I provide examples of research into diseases of threatened species that have applied comparative molecular approaches.

Comparative genomics of wildlife can support the identification of candidate loci responsible for heritable disorders that may increase in frequency in small populations as a result of inbreeding. An example is the lethal chondrodystrophy seen in the critically endangered California condor *Gymnogyps californianus* [16]. This disease showed Mendelian segregation in condor pedigrees [62], so researchers developed genomic resources with the aim of identifying the genetic basis of the disorder and revealing genetic markers linked to the disease [16,63]. It is hoped that this ongoing research will ultimately inform the design of captive breeding protocols that can reduce the frequency of chondrodystrophy, although it would be important to avoid losses of genetic diversity overall (i.e. at other loci), and ensure inbreeding is minimised to avoid further decreases in individual fitness [23].

A particularly interesting application of genomics to the study of disease emergence in a threatened species is in the Tasmanian devil *Sarcophilus harrisi*, which is under threat from devil facial tumour disease (DFTD). DFTD is a highly contagious and fatal transmissible cancer that has caused dramatic population declines in the species' native Tasmania, since its first observation in 1996 [64]. Uniquely, DFTD cells are the infectious agent of the disease, spread via direct contact between individuals [65], probably when devils bite one another during aggressive interactions [66]. In this case, comparative genomic research compares the host genome with that of its “pathogen”, DFTD, via whole-genome sequencing [67]. The aims of this work are to identify the underlying mutations driving tumorigenesis and the ability of DFTD to evade the immune system of the host, with a long-term view toward a potentially identifying a target for a vaccine or other therapeutic [67, 68]. Many insights into the origin, transmission and evolution of DFTD have arisen from this and related work, which is still ongoing [68]. Meanwhile, comparisons between the genome of DFTD and its host have enabled the identification of marker loci, which, through genotyping of large numbers of tumour samples from throughout the population, have confirmed the clonal origin of DFTD: it only arose once and has spread [67]. This result has had important conservation implications via the realisation that healthy devils could be prevented from acquiring the disease if they were physically separated from diseased animals – leading to the establishment of a highly successful captive “insurance” population [69].

In addition to these examples of particular disease processes, comparative genomics can be a powerful tool for the study of a species' immunome. By examining the diverse characteristics of species' immunity genes, we can learn more about how each species has evolved to respond to pathogens in its environment [70]. In a conservation context, the comparative approach can be used to address questions about the particular characteristics that predispose populations to emerging threats (as has been done from an ecological perspective [9]). For example, recent world-wide amphibian declines have resulted from the spread of the chytrid fungus *Batrachochytrium dendrobatidis* [71]. Comparative immunogenomics of *Xenopus* and closely related species allows researchers to examine the response of the model amphibians to chytrid and other pathogens, in order to discover how diseases may affect threatened species [72]. As whole-genome sequences are

published for a greater number of species, the hope is that whole-immunome comparisons across species will become more refined, improving the power of studies examining the immunogenetic diversity, and therefore the disease consequences of low diversity, in threatened taxa [18]. At present, immune-gene regions are challenging to assemble, in part due to the very traits that make immunity highly adaptable: high levels of gene duplication [73,74]. It is therefore difficult to determine how many copies of duplicate genes are present in a species or even individual genome [75,76]. Emerging technologies offering longer sequencing reads [e.g. 77], as well as continued development and assessment of computational approaches [e.g. 78,79], may help to overcome this challenge.

2. Conclusions

I have presented a brief overview of conservation genetics questions that can be targeted by a comparative genomic approach: including the relationships between related species or populations (delineation of conservation units and studies of hybridisation), studies of cross-species variation (examination of the genomic regions contributing to species distinctiveness and divergence) and the interactions between species and their threatening processes (whether intrinsic or extrinsic). It is clear that many aspects of preserving threatened species diversity can benefit from looking beyond the threatened species itself, and considering the differences between the rare species' genome and other genomes that share its environment, although empirical examples in this research area, utilising threatened species themselves, are scarce. A particularly exciting path of inquiry is the application of comparative genomics to the study of the ecological context of genomes, through a better understanding of interactions between species, such as hosts and parasites, in an “extended phenotype”, co-evolutionary framework [80,81] (for examples see [82–84]). Such findings present promising opportunities to better understand the evolutionary consequences of anthropogenic biotic disturbances to ecosystems, such as the shifting distributions of pathogens, predators, prey and competitors. A better understanding of these complex ecological interactions can help us to understand the relationships amongst species and inform conservation planning at higher levels, such as the protection of vulnerable habitats and ecological processes. The preservation of biodiversity is essential at multiple levels: from molecular diversity, to species, to whole ecosystems and habitats [e.g. 85]. One of the best arguments for the use of comparative genomics in conservation is that whole-genome level data is necessary to monitor and protect the greatest breadth of genetic biodiversity [1]. By comparing and contrasting the processes that influence the genomic composition of threatened species with those of more-common species, we can identify what makes vulnerable species ecologically and genetically unique, and stand the best chance at preserving them and their individual roles in the global ecosystem.

Acknowledgments

This manuscript was improved by suggestions from three anonymous reviewers. I thank K. Belov for valuable conversations on this topic. I acknowledge the support of Australian Research Council, San Diego Zoo Global, Save the Tasmanian Devil Program, Zoo and Aquarium Association and University of Sydney.

References

- [1] McMahon BJ, Teeling EC, Höglund J. How and why should we implement genomics into conservation? *Evol App* 2014;7:999–1007.
- [2] Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, Diekhans M, et al. The UCSC Genome Browser database: 2014 update. *Nucleic Acids Res* 2014;42:D764–70.
- [3] Ellegren H. Genome sequencing and population genomics in non-model organisms. *Trends Ecol Evol* 2014;29:51–63.
- [4] Kohn MH, Murphy WJ, Ostrander EA, Wayne RK. Genomics and conservation genetics. *Trends Ecol Evol* 2006;21:629–37.

- [5] Luikart G, England PR, Tallmon D, Jordan S, Taberlet P. The power and promise of population genomics: from genotyping to genome typing. *Nat Rev Genet* 2003;4: 981–94.
- [6] Shafer ABA, Wolf JBW, Alves PC, Bergström L, Bruford MW, Brännström I, et al. Genomics and the challenging translation into conservation practice. *Trends Ecol Evol* 2015;30:78–87.
- [7] Frankham R. Conservation genetics. *Annu Rev Genet* 1995;29:305–27.
- [8] Purvis A, Gittleman JL, Cowlishaw G, Mace GM. Predicting extinction risk in declining species. *Proc Roy Soc Lond B: Biol Sci* 2000;267:1947–52.
- [9] Fisher DO, Owens IPF. The comparative method in conservation biology. *Trends Ecol Evol* 2004;19:391–8.
- [10] Twyford AD, Ennos RA. Next-generation hybridization and introgression. *Heredity* 2012;108:179–89.
- [11] Davey JW, Hohenlohe PA, Etter PD, Boone JQ, Catchen JM, Blaxter ML, et al. Genome-wide genetic marker discovery and genotyping using next-generation sequencing. *Nat Rev Genet* 2011;12:499–510.
- [12] McCormack JE, Hird SM, Zellmer AJ, Carstens BC, Brumfield RT. Applications of next-generation sequencing to phylogeography and phylogenetics. *Mol Phylogenet Evol* 2013;66:526–38.
- [13] Perry GH, Melsted P, Marioni JC, Wang Y, Bainer R, Pickrell JK, et al. Comparative RNA sequencing reveals substantial genetic variation in endangered primates. *Genom Res* 2012;22:602–10.
- [14] vonHoldt BM, Pollinger JP, Earl DA, Knowles JC, Boyko AR, Parker H, et al. A genome-wide perspective on the evolutionary history of enigmatic wolf-like canids. *Genom Res* 2011;21:1294–305.
- [15] Miller JM, Kijas JW, Heaton MP, McEwan JC, Coltman DW. Consistent divergence times and allele sharing measured from cross-species application of SNP chips developed for three domestic species. *Mol Ecol* 2012;21:1145–50.
- [16] Romanov M, Tuttle E, Houck M, Modi W, Chemnick L, Korody M, et al. The value of avian genomics to the conservation of wildlife. *BMC Genom* 2009;10:S10.
- [17] Toro MA, Villanueva B, Fernández J. Genomics applied to management strategies in conservation programmes. *Livest Sci* 2014;166:48–53.
- [18] Larsen PA, Campbell CR, Yoder AD. Next-generation approaches to advancing ecomonogenomic research in critically endangered primates. *Mol Ecol Resour* 2014;14:1198–209.
- [19] Locke DP, Hillier LW, Warren WC, Worley KC, Nazareth LV, Muzny DM, et al. Comparative and demographic analysis of orang-utan genomes. *Nature* 2011;469:529–33.
- [20] Hohenlohe PA, Amish SJ, Catchen JM, Allendorf FW, Luikart G. Next-generation RAD sequencing identifies thousands of SNPs for assessing hybridization between rainbow and westslope cutthroat trout. *Mol Ecol Resour* 2011;11:117–22.
- [21] Steiner CC, Putnam AS, Hoek PEA, Ryder OA. Conservation genomics of threatened animal species. *Annu Rev Anim Biosci* 2013;1:261–81.
- [22] Alföldi J, Lindblad-Toh K. Comparative genomics as a tool to understand evolution and disease. *Genom Res* 2013;23:1063–8.
- [23] Frankham R. Challenges and opportunities of genetic approaches to biological conservation. *Biol Conserv* 2010;143:1919–27.
- [24] Palsboll PJ, Bérubé M, Allendorf FW. Identification of management units using population genetic data. *Trends Ecol Evol* 2007;22:11–6.
- [25] Moritz C. Defining 'evolutionarily significant units' for conservation. *Trends Ecol Evol* 1994;9:373–5.
- [26] Gebremedhin B, Ficetola GF, Naderi S, Rezaei HR, Maudet C, Rioux D, et al. Frontiers in identifying conservation units: from neutral markers to adaptive genetic variation. *Anim Conserv* 2009;12:107–9.
- [27] Piertney SB, Oliver MK. The evolutionary ecology of the major histocompatibility complex. *Heredity* 2006;96:7–21.
- [28] Zhu Y, Wan Q-H, Yu B, Ge Y-F, Fang S-G. Patterns of genetic differentiation at MHC class I genes and microsatellites identify conservation units in the giant panda. *BMC Evol Biol* 2013;13:227.
- [29] Vásquez-Carrillo C, Friesen V, Hall L, Peery MZ. Variation in MHC class II B genes in marbled murrelets: implications for delineating conservation units. *Anim Conserv* 2014;17:244–55.
- [30] Acevedo-Whitehouse K, Cunningham AA. Is MHC enough for understanding wildlife immunogenetics? *Trends Ecol Evol* 2006;21:433–8.
- [31] Wagner CE, Keller I, Wittwer S, Selz OM, Mwaiko S, Greuter L, et al. Genome-wide RAD sequence data provide unprecedented resolution of species boundaries and relationships in the Lake Victoria cichlid adaptive radiation. *Mol Ecol* 2013;22:787–98.
- [32] Isaac NJB, Turvey ST, Collen B, Waterman C, Baillie JEM. Mammals on the EDGE: conservation priorities based on threat and phylogeny. *PLoS One* 2007;2:e296.
- [33] Funk WC, McKay JK, Hohenlohe PA, Allendorf FW. Harnessing genomics for delineating conservation units. *Trends Ecol Evol* 2012;27:489–96.
- [34] Anderson CNK, Liu L, Pearl D, Edwards SV. Tangled trees: the challenge of inferring species trees from coalescent and noncoalescent genes. In: Anisimova M, editor. *Evolutionary genomics*. Humana Press; 2012. p. 3–28.
- [35] Earl D, Nguyen N, Hickey G, Harris RS, Fitzgerald S, Beal K, et al. Alignathon: a competitive assessment of whole-genome alignment methods. *Genom Res* 2014;24: 2077–89.
- [36] Sutton JT, Nakagawa S, Robertson BC, Jamieson IG. Disentangling the roles of natural selection and genetic drift in shaping variation at MHC immunity genes. *Mol Ecol* 2011;20:4408–20.
- [37] Steeves TE, Maloney RF, Hale ML, Tylianakis JM, Gemmill NJ. Genetic analyses reveal hybridization but no hybrid swarm in one of the world's rarest birds. *Mol Ecol* 2010; 19:5090–100.
- [38] Rhymer JM, Simberloff D. Extinction by hybridization and introgression. *Annu Rev Ecol Syst* 1996;27:83–109.
- [39] Allendorf FW, Leary RF, Spruell P, Wenburg JK. The problems with hybrids: setting conservation guidelines. *Trends Ecol Evol* 2001;16:613–22.
- [40] Frankham R, Ballou JD, Eldridge MDB, Lacy RC, Ralls K, Dudash MR, et al. Predicting the probability of outbreeding depression. *Conserv Biol* 2011;25:465–75.
- [41] Bourret V, O'Reilly PT, Carr JW, Berg PR, Bernatchez L. Temporal change in genetic integrity suggests loss of local adaptation in a wild Atlantic salmon (*Salmo salar*) population following introgression by farmed escapees. *Heredity* 2011;106:500–10.
- [42] Muhlfeld CC, Kalinowski ST, McMahon TE, Taper ML, Painter S, Leary RF, et al. Hybridization rapidly reduces fitness of a native trout in the wild. *Biol Lett* 2009;5: 328–31.
- [43] Hedrick PW. Conservation genetics: where are we now? *Trends Ecol Evol* 2001;16: 629–36.
- [44] Van Doornik DM, Eddy DL, Waples RS, Boe SJ, Hoffnagle TL, Berntson EA, et al. Genetic monitoring of threatened Chinook salmon populations: estimating introgression of nonnative hatchery stocks and temporal genetic changes. *N Am J Fish Manage* 2013;33:693–706.
- [45] Tiwari V, Wang S, Sehgal S, Vrana J, Friebe B, Kubalaková M, et al. SNP discovery for mapping alien introgressions in wheat. *BMC Genomics* 2014;15:273.
- [46] Hohenlohe PA, Day MD, Amish SJ, Miller MR, Kamps-Hughes N, et al. Genomic patterns of introgression in rainbow and westslope cutthroat trout illuminated by overlapping paired-end RAD sequencing. *Mol Ecol* 2013;22:3002–13.
- [47] Shepard BB, May BE, Urie W. Status and conservation of westslope cutthroat trout within the western United States. *N Am J Fish Manage* 2005;25:1426–40.
- [48] Amador C, Toro MÁ, Fernández J. Molecular markers allow to remove introgressed genetic background: a simulation study. *PLoS One* 2012;7:e49409.
- [49] Amador C, Hayes BJ, Daetwyler HD. Genomic selection for recovery of original genetic background from hybrids of endangered and common breeds. *Evol Appl* 2014;7: 227–37.
- [50] Amador C, Toro M, Fernández J. Removing exogenous information using pedigree data. *Conserv Genet* 2011;12:1565–73.
- [51] Simčič M, Smetko A, Sölkner J, Seichter D, Gorjanc G, Kompan D, et al. Recovery of native genetic background in admixed populations using haplotypes, phenotypes, and pedigree information—using Cika cattle as a case breed. *PLoS One* 2015;10: e0123253.
- [52] Zhan X, Pan S, Wang J, Dixon A, He J, Muller MG, et al. Peregrine and saker falcon genome sequences provide insights into evolution of a predatory lifestyle. *Nat Genet* 2013;45:563–6.
- [53] Montgomery ME, Woodworth LM, England PR, Briscoe DA, Frankham R. Widespread selective sweeps affecting microsatellites in *Drosophila* populations adapting to captivity: implications for captive breeding programs. *Biol Conserv* 2010;143:1842–9.
- [54] Mikkelsen TS, Wakefield MJ, Aken B, Amemiya CT, Chang JL, Duke S, et al. Genome of the marsupial *Monodelphis domestica* reveals innovation in non-coding sequences. *Nature* 2007;447:167–77.
- [55] Li R, Fan W, Tian G, Zhu H, He L, Cai J, et al. The sequence and de novo assembly of the giant panda genome. *Nature* 2010;463:311–7.
- [56] Nei M, Rooney AP. Concerted and birth-and-death evolution of multigene families. *Annu Rev Genet* 2005;39:121–52.
- [57] Denton JF, Lugo-Martinez J, Tucker AE, Schrider DR, Warren WC, Hahn MW. Extensive error in the number of genes inferred from draft genome assemblies. *PLoS One* 2014;10:e1003998.
- [58] Ekblom R, Wolf JBW. A field guide to whole-genome sequencing, assembly and annotation. *Evol App* 2014;7:1026–42.
- [59] Pavlidis P, Jensen JD, Stephan W, Stamatakis A. A critical assessment of storytelling: gene ontology categories and the importance of validating genomic scans. *Mol Biol Evol* 2012;29:3237–48.
- [60] Spielman D, Brook BW, Briscoe DA, Frankham R. Does inbreeding and loss of genetic diversity decrease disease resistance? *Conserv Genet* 2004;5:439–48.
- [61] Frankham R, Ballou JD, Briscoe DA. Introduction to conservation genetics. Cambridge: Cambridge University Press; 2002;640.
- [62] Ralls K, Ballou JD, Rideout BA, Frankham R. Genetic management of chondrodystrophy in California condors. *Anim Conserv* 2000;3:145–53.
- [63] Walters JR, Derrickson SR, Fry DM, Haig SM, Marzluff JM, Wunderle JMJ. Status of the California condor (*Gymnogyps californianus*) and efforts to achieve its recovery. *Auk* 2010;127:969–1001.
- [64] Hawkins CE, Baars C, Hesterman H, Hocking GJ, Jones ME, Lazenby B, et al. Emerging disease and population decline of an island endemic, the Tasmanian devil *Sarcophilus harrisii*. *Biol Conserv* 2006;131:307–24.
- [65] Pearce A, Swift K. Allograft theory: transmission of devil facial-tumour disease. *Nature* 2006;439:549.
- [66] Hamede RK, McCallum H, Jones M. Biting injuries and transmission of Tasmanian devil facial tumour disease. *J Anim Ecol* 2013;82:182–90.
- [67] Murchison EP, Schulz-Trieglaff OB, Ning Z, Alexandrov LB, Bauer MJ, Fu B, et al. Genome sequencing and analysis of the Tasmanian devil and its transmissible cancer. *Cell* 2012;148:780–91.
- [68] Deakin JE, Belov K. A comparative genomics approach to understanding transmissible cancer in Tasmanian devils. *Annu Rev Genomics Hum Genet* 2012; 13:207–22.
- [69] DPIPWE. Save the Tasmanian Devil Program 2012–13 annual program report. Hobart: Department of Primary Industries, Parks, Water and Environment; 2013.
- [70] Criscitiello MF, de Figueiredo P. Fifty shades of immune defense. *PLoS Pathog* 2013; 9:e1003110.
- [71] Lips KR, Diffendorfer J, Mendelson III JR, Sears MW. Riding the wave: reconciling the roles of disease and climate change in amphibian declines. *PLoS Biol* 2008;6:e72.
- [72] Robert J, Ohta Y. Comparative and developmental study of the immune system in *Xenopus*. *Dev Dyn* 2009;238:1249–70.
- [73] Nei M, Gu X, Sitnikova T. Evolution by the birth-and-death process in multigene families of the vertebrate immune system. *Proc Nat Acad Sci USA* 1997;94: 7799–806.

- [74] Temperley N, Berlin S, Paton I, Griffin D, Burt D. Evolution of the chicken Toll-like receptor gene family: a story of gene gain and gene loss. *BMC Genom* 2008;9.
- [75] Cheng Y, Stuart A, Morris K, Taylor R, Siddle H. Antigen-presenting genes and genomic copy number variations in the Tasmanian devil MHC. *BMC Genom* 2012;13:87.
- [76] Alcaide M, Muñoz J, Martínez-de la Puente J, Soriguer R, Figuerola J. Extraordinary MHC class II B diversity in a non-passerine, wild bird: the Eurasian Coot *Fulica atra* (Aves: Rallidae). *Ecol Evol* 2014;4:688–98.
- [77] English AC, Richards S, Han Y, Wang M, Vee V, Qu J, et al. Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology. *PLoS One* 2012;7:e47768.
- [78] Salzberg SL, Phillippy AM, Zimin A, Puiu D, Magoc T, Koren S, et al. GAGE: a critical evaluation of genome assemblies and assembly algorithms. *Genome Res* 2012;22:557–67.
- [79] Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 2013;10:563–9.
- [80] Dawkins R. *The extended phenotype*. Oxford: Oxford University Press; 1982.
- [81] Dawkins R. *The extended phenotype: the long reach of the gene*. Oxford: Oxford University Press; 1999.
- [82] Lambrechts L, Fellous S, Koella JC. Coevolutionary interactions between host and parasite genotypes. *Trends Parasitol* 2006;22:12–6.
- [83] Foth BJ, Tsai IJ, Reid AJ, Bancroft AJ, Nichol S, Tracey A, et al. Whipworm genome and dual-species transcriptome analyses provide molecular insights into an intimate host–parasite interaction. *Nat Genet* 2014;46:693–700.
- [84] Pilosof S, Fortuna MA, Cosson JF, Galan M, Kittipong C, Ribas A, et al. Host–parasite network structure is associated with community-level immunogenetic diversity. *Nat Commun* 2014;5:5172.
- [85] Franklin JF. Preserving biodiversity: species, ecosystems, or landscapes? *Ecol Appl* 1993;3:202–5.